Author Index

Allen, E., Myers, C., Frank, D., Pettigrew, A., Thompson, S. and Blumer, J. L.
Application of an organic solvent extraction to the determination of catechol-O-methyltransferase activity by high-performance liquid chromatography in human mononuclear cells 578(1992)175

Al Odaib, A., see Leis, H. J. 578(1992)116

Alonso, R. M., see Rodríguez, F. J. 578(1992)146

Alvarez-Bujidos, M. L., Ortiz, A., Balaña, R., Cubría, J. C., Ordoñez, D. and Negro, A.
Determination of luxabendazole in biological fluids by high-performance liquid chromatography 578(1992)321

Alvaro, R. F., Rosegay, A. and Chiu, S.-H. L.
Determination of enantiomeric concentrations of a 2,5-diaryltetrahydrofuran (L-668,750), a platelet-activating factor receptor antagonist, in rat plasma using a chiral α_1-acid glycoprotein high-performance liquid chromatographic column 578(1992)327

Anderson, R. E., see Chen, H. 578(1992)124

André, D., see Vérité, P. 578(1992)134

Bailey, L. C., see Wilhelm, J. A. 578(1992)231

Balaña, R., see Alvarez-Bujidos, M. L. 578(1992)321

Bell, C. J., see Bernert, Jr., J. T. 578(1992)1

Bernert, Jr., J. T., Bell, C. J., McGuffey, J. E. and Waymack, P. P.
Determination of "free" glycerol in human serum reference materials by isotope-dilution gas chromatography–mass spectrometry 578(1992)1

Blagbrough, I. S., Daykin, M. M., Doherty, M., Pattrick, M. and Shaw, P. N.
High-performance liquid chromatographic determination of naproxen, ibuprofen and diclofenac in plasma and synovial fluid in man 578(1992)251

Blaschke, G., see Westhoff, F. 578(1992)265

Blumer, J. L., see Allen, E. 578(1992)175

Boje, K. M., see Miller, R. 578(1992)103

Breda, M., Pianezzola, E. and Strolin Benedetti, M.
Determination of 3′-deamino-3′-[2(*S*)-methoxy-4-morpholinyl]doxorubicin, a new morpholinyl anthracycline, in plasma by high-performance liquid chromatography with fluorescence detection 578(1992)309

Brown, A. R., James, C. D., Welch, R. M. and Harrelson, J. C.
Stereoselective high-performance liquid chromatographic assay with fluorometric detection for the isomers of mivacurium in human plasma 578(1992)302

Carlqvist, J., see Lanbeck-Vallén, K. 578(1992)71

Chen, H. and Anderson, R. E.
Quantitation of phenacyl esters of retinal fatty acids by high-performance liquid chromatography 578(1992)124

Chiu, S.-H. L., see Alvaro, R. F. 578(1992)327

Chovan, J. P., Zane, P. A. and Greenberg, G. E.
Automated-extraction, high-performance liquid chromatographic method and pharmacokinetics in rats of a highly A_2-selective adenosine agonist, CGS 21680 578(1992)77

Christensen, J. M., see Hansen, Å. M. 578(1992)85

Cissik, J. H., see Schwertner, H. A. 578(1992)297

Cubría, J. C., see Alvarez-Bujidos, M. L. 578(1992)321

Dakshinamurti, K., see Sharma, S. K. 578(1992)45

Daykin, M. M., see Blagbrough, I. S. 578(1992)251

De Neve, W., see Van Belle, S. J.-P. 578(1992)223

De Smet, M., see Van Belle, S. J.-P. 578(1992)223

Deyl, Z., see Hodny, Z. 578(1992)53

Doherty, M., see Blagbrough, I. S. 578(1992)251

Dostert, P., see Niwa, T. 578(1992)109

Etoh, T., Miyazaki, M., Harada, K., Nakayama, M. and Sugii, A.
Rapid analysis of human serum albumin by high-performance liquid chromatography 578(1992)292

Evans, R. R. and Relling, M. V.
Automated high-performance liquid chromatographic assay for the determination of 7-ethoxycoumarin and umbelliferone 578(1992)141

Frank, D., see Allen, E. 578(1992)175

Fried, K., see Lloyd, D. K. 578(1992)283

Fürst, W. and Hallström, S.
Simultaneous determination of myocardial nucleotides, nucleosides, purine bases and creatine phosphate by ion-pair high-performance liquid chromatography 578(1992)39

Gallo, J. M., see Qian, M. 578(1992)316

Gleispach, H., see Leis, H. J. 578(1992)116

Goto, N., Kamata, T. and Ikegami, K.
Trace analysis of quinapril and its active metabolite, quinaprilat, in human plasma and urine by gas chromatography–negative-ion chemical ionization mass spectrometry 578(1992)195

Goto, N., Sato, T., Shigetoshi, M. and Ikegami, K.
Determination of dioxopiperazine metabolites of quinapril in biological fluids by gas chromatography–mass spectrometry 578(1992)203

Greenberg, G. E., see Chovan, J. P. 578(1992)77

Gulati, A.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection. Reply to Zweipfenning 578(1992)335

Hallström, S., see Fürst, W. 578(1992)39

Hansen, Å. M., Poulsen, O. M., Christensen, J. M. and Hansen, S. H.
Determination of 2-naphthylamine in urine by a novel reversed-phase high-performance liquid chromatography method 578(1992)85

Hansen, S. H., see Hansen, Å. M. 578(1992)85

Harada, K., see Etoh, T. 578(1992)292

Harrelson, J. C., see Brown, A. R. 578(1992)302

Härtter, S. and Hiemke, C.
Column switching and high-performance liquid chromatography in the analysis of amitriptyline, nortriptyline and hydroxylated metabolites in human plasma or serum 578(1992)273

Hiemke, C., see Härtter, S. 578(1992)273
Hodny, Z., Struzinsky, R. and Deyl, Z.
Silver staining of collagen type I after sodium dodecylsulphate polyacrylamide gel electrophoresis: effect of Maillard reaction 578(1992)53
Horváth, G., see Tomori, É. 578(1992)91
Hsieh, J. Y.-K., see Kline, W. F. 578(1992)31
Ikegami, K., see Goto, N. 578(1992)195
Ikegami, K., see Goto, N. 578(1992)203
Ikegami, Y., see Sano, K. 578(1992)63
James, C. D., see Brown, A. R. 578(1992)302
Jiménez, R. M., see Rodríguez, F. J. 578(1992)146
Jongen, M. J. M., see Leenheers, L. H. 578(1992)189
Kalasinsky, K. S., see Kuhlman, Jr., J. J. 578(1992)207
Kamata, T., see Goto, N. 578(1992)195
Kataoka, H., Ohrui, S., Miyamoto, Y. and Makita, M.
Determination of urinary β-phenylethylamine as its N-benzenesulphonamide derivative by gas chromatography with flame photometric detection 578(1992)120
Kinosita, I., see Sano, K. 578(1992)63
Kinter, M., Sullivan, S., Roberts, R. J. and Spitz, D.
Trace quantitation of 4-hydroxy-2-nonenal in biological samples as its oxime–bis-*tert*.-butyldimethylsilyl derivative using 3-hydroxynonanal as an internal standard 578(1992)9
Kissinger, P. T., see Linhares, M. C. 578(1992)157
Klette, K. L., see Kuhlman, Jr., J. J. 578(1992)207
Kline, W. F., Matuszewski, B. K. and Hsieh, J. Y.-K.
Determination of a cyclic heptapeptide, a novel fibrinogen receptor antagonist, in human plasma by high-performance liquid chromatography with automated pre-column derivatization, column switching and fluorescence detection 578(1992)31
Kuhlman, Jr., J. J., Levine, B., Klette, K. L., Magluilo, Jr., J., Kalasinsky, K. S. and Smith, M. L.
Measurement of azacyclonol in urine and serum of humans following terfenadine (Seldane) administration using gas chromatography–mass spectrometry 578(1992)207
Kurono, M., Yoshida, K. and Naruto, S.
Determination of AJ-3941, a possible agent for the treatment of cerebrovascular disorders, in plasma and brain by means of high-performance liquid chromatography with fluorescence detection 578(1992)152
Lafont, O., see Vérité, P. 578(1992)134
La Grone, J., see Miller, R. 578(1992)103
Lanbeck-Vallén, K. and Carlqvist, J.
Determination of cefcanel in plasma and urine by high-performance liquid chromatography using coupled columns, after administration of the new cephalosporin prodrug cefcanel daloxate hydrochloride 578(1992)71
Lee Chin, J. R.
Determination of the catecholamines and serotonin, their precursors tyrosine and tryptophan, and their main metabolites in rat brain using reversed-phase high-performance liquid chromatography with fluorimetric and oxidative amperometric detection in series 578(1992)17
Leenheers, L. H., Van Breugel, D. C., Ravensberg, J. C., Meuling, W. J. A. and Jongen, M. J. M.
Determination of 2-isopropoxyphenol in urine by capillary gas chromatography and mass-selective detection 578(1992)189
Leis, H. J., Ozand, P. T., Al Odaib, A. and Gleispach, H.
Quantitative measurement of N-acetyl-L-aspartic acid in urine by gas chromatography with negative-ion chemical ionization mass spectrometry 578(1992)116
Levine, B., see Kuhlman, Jr., J. J. 578(1992)207
Linhares, M. C. and Kissinger, P. T.
In vivo sampling using loop microdialysis probes coupled to a liquid chromatograph 578(1992)157
Linhart, I., Vosmanská, M., Mitera, J. and Šmejkal, J.
Biotransformation of diethenylbenzenes. IV. A simple high-performance liquid chromatographic method for separation of urinary metabolites of 1,3-diethenylbenzene on analytical and semi-preparative scales 578(1992)215
Lloyd, D. K., Fried, K. and Wainer, I. W.
Determination of N-acetylator phenotype using caffeine as a probe compound: a comparison of high-performance liquid chromatography and capillary electrophoresis methods 578(1992)283
Magluilo, Jr., J., see Kuhlman, Jr., J. J. 578(1992)207
Makita, M., see Kataoka, H. 578(1992)120
Maruyama, W., see Niwa, T. 578(1992)109
Massart, D. L., see Van Belle, S. J.-P. 578(1992)223
Matuszewski, B. K., see Kline, W. F. 578(1992)31
McGuffey, J. E., see Bernert, Jr., J. T. 578(1992)1
Medzihradszky, D., see Urbányi, Z. 578(1992)130
Ménager, S., see Vérité, P. 578(1992)134
Mészáros, S., see Tomori, É. 578(1992)91
Meuling, W. J. A., see Leenheers, L. H. 578(1992)189
Mihara, R., see Sano, K. 578(1992)63
Miller, R., La Grone, J., Skolnick, P. and Boje, K. M.
High-performance liquid chromatographic assay for 1-aminocyclopropanecarboxylic acid from plasma and brain 578(1992)103
Mitera, J., see Linhart, I. 578(1992)215
Miyamoto, Y., see Kataoka, H. 578(1992)120
Miyazaki, M., see Etoh, T. 578(1992)292
Monsaert, C., see Van Belle, S. J.-P. 578(1992)223
Myers, C., see Allen, E. 578(1992)175
Nagatsu, T., see Niwa, T. 578(1992)109
Nakahara, D., see Niwa, T. 578(1992)109
Nakayama, M., see Etoh, T. 578(1992)292
Naoi, M., see Niwa, T. 578(1992)109
Naruto, S., see Kurono, M. 578(1992)152
Negro, A., see Alvarez-Bujidos, M. L. 578(1992)321
Niwa, T., Maruyama, W., Nakahara, D., Takeda, N., Yoshizumi, H., Tatematsu, A., Takahashi, A., Dostert, P., Naoi, M. and Nagatsu, T.
Endogenous synthesis of N-methylsalsolinol, an analogue of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, in rat brain during *in vivo* microdialysis with salsolinol, as demonstrated by gas chromatography–mass spectrometry 578(1992)109
Ohrui, S., see Kataoka, H. 578(1992)120
Ordoñez, D., see Alvarez-Bujidos, M. L. 578(1992)321
Ortiz, A., see Alvarez-Bujidos, M. L. 578(1992)321
Ozand, P. T., see Leis, H. J. 578(1992)116
Pátfalusi, M., see Tomori, É. 578(1992)91

Patterson, W. R., see Schwertner, H. A. 578(1992)297
Pattrick, M., see Blagbrough, I. S. 578(1992)251
Pettigrew, A., see Allen, E. 578(1992)175
Pianezzola, E., see Breda, M. 578(1992)309
Platt, D., see Rieck, W. 578(1992)259
Poulsen, O. M., see Hansen, Å. M. 578(1992)85
Qian, M. and Gallo, J. M.
High-performance liquid chromatographic determination of the calcium channel blocker nimodipine in monkey plasma 578(1992)316
Ravensberg, J. C., see Leenheers, L. H. 578(1992)189
Relling, M. V., see Evans, R. R. 578(1992)141
Rieck, W. and Platt, D.
High-performance liquid chromatographic method for the determination of alprazolam in plasma using the column-switching technique 578(1992)259
Roberts, R. J., see Kinter, M. 578(1992)9
Rodríguez, F. J., Jiménez, R. M. and Alonso, R. M.
Separation and determination of aminohalogenbenzophenones by high-performance liquid chromatography with electrochemical detection 578(1992)146
Rosegay, A., see Alvaro, R. F. 578(1992)327
Sano, K., Kinosita, I., Mihara, R., Ikegami, Y. and Uesugi, T.
High-performance liquid chromatographic determination of sulphobromophthalein and its conjugates 578(1992)63
Sato, T., see Goto, N. 578(1992)203
Schwertner, H. A., Patterson, W. R., Cissik, J. H. and Wilson, K. W.
New extraction procedure and high-performance liquid chromatographic method for analyzing polyethylene glycol-400 in urine 578(1992)297
Sharma, S. K. and Dakshinamurti, K.
Determination of vitamin B_6 vitamers and pyridoxic acid in biological samples 578(1992)45
Shaw, P. N., see Blagbrough, I. S. 578(1992)251
Shepard, T. A., see Wilhelm, J. A. 578(1992)231
Shigetoshi, M., see Goto, N. 578(1992)203
Skolnick, P., see Miller, R. 578(1992)103
Šmejkal, J., see Linhart, I. 578(1992)215
Smith, M. L., see Kuhlman, Jr., J. J. 578(1992)207
Spitz, D., see Kinter, M. 578(1992)9
Storme, G. A., see Van Belle, S. J.-P. 578(1992)223
Strolin Benedetti, M., see Breda, M. 578(1992)309
Struzinsky, R., see Hodny, Z. 578(1992)53
Sugii, A., see Etoh, T. 578(1992)292
Sullivan, S., see Kinter, M. 578(1992)9
Takahashi, A., see Niwa, T. 578(1992)109
Takeda, N., see Niwa, T. 578(1992)109
Tatematsu, A., see Niwa, T. 578(1992)109
Thompson, S., see Allen, E. 578(1992)175
Tomori, É., Horváth, G., Pátfalusi, M., Mészáros, S. and Vereczkey, L.
Pharmacokinetic and metabolism studies on girisopam by chromatographic and spectrometric methods in humans 578(1992)91
Uesugi, T., see Sano, K. 578(1992)63
Urbányi, Z. and Medzihradszky, D.
Rapid method to isolate serum amyloid P component from human plasma. Characterization of the isolated protein 578(1992)130
Van Belle, S. J.-P., De Smet, M., De Neve, W., Monsaert, C., Storme, G. A. and Massart, D. L.
Determination of vinca alkaloids in mouse tissues by high-performance liquid chromatography 578(1992)223
Van Breugel, D. C., see Leenheers, L. H. 578(1992)189
Van den Biggelaar-Martea, M., see Vree, T. B. 578(1992)239
Venturella, V. S., see Wilhelm, J. A. 578(1992)231
Vereczkey, L., see Tomori, É. 578(1992)91
Vérité, P., André, D., Ménager, S. and Lafont, O.
Gas chromatographic–mass spectrometric method for analysis of a pyrimido-*s*-triazine and some of its metabolites in dog urine 578(1992)134
Verwey-van Wissen, C. P. W. G. M., see Vree, T. B. 578(1992)239
Volin, P.
Determination of urinary normetanephrine, metanephrine and 3-methoxytyramine by high-performance liquid chromatography with electrochemical detection: comparison between automated column-switching and manual dual-column sample purification methods 578(1992)165
Vosmanská, M., see Linhart, I. 578(1992)215
Vree, T. B., Van den Biggelaar-Martea, M. and Verwey-van Wissen, C. P. W. G. M.
Determination of naproxen and its metabolite O-desmethylnaproxen with their acyl glucuronides in human plasma and urine by means of direct gradient high-performance liquid chromatography 578(1992)239
Wainer, I. W., see Lloyd, D. K. 578(1992)283
Waymack, P. P., see Bernert, Jr., J. T. 578(1992)1
Welch, R. M., see Brown, A. R. 578(1992)302
Westhoff, F. and Blaschke, G.
High-performance liquid chromatographic determination of the stereoselective biotransformation of the chiral drug praziquantel 578(1992)265
Wilhelm, J. A., Bailey, L. C., Shepard, T. A. and Venturella, V. S.
Simultaneous determination of phenolphthalein and phenolphthalein glucuronide from dog serum, urine and bile by high-performance liquid chromatography 578(1992)231
Wilson, K. W., see Schwertner, H. A. 578(1992)297
Yoshida, K., see Kurono, M. 578(1992)152
Yoshizumi, H., see Niwa, T. 578(1992)109
Zane, P. A., see Chovan, J. P. 578(1992)77
Zweipfenning, P. G. M.
High-performance liquid chromatographic determination of morphine, morphine-3-glucuronide, morphine-6-glucuronide and codeine in biological samples using multi-wavelength forward optical detection: a reply 578(1992)333